In re Application of: Khan, Saeed R. Application No.: 10/596,751 Filed: October 18, 2007

Page 2

Amendments to the Claims:

Please amend claims 1-9 as shown in the listing of the claims.

Please cancel claim 10 without prejudice.

Please add claims 17-19 as shown in the listing of the claims.

This listing of the claims will replace all prior listing of the claims in this application.

Listing of the Claims

(Currently Amended) A compound having the formula:

wherein

$$X^1$$
 is $-\Theta$ $C = \text{ or } -N =$:

 ${\bf L}^1$ is a bond, substituted or unsubstituted alkylene, or substituted or unsubstituted heteroalkylene; and

R1 has the formula:

$$\underbrace{\frac{\xi}{\xi} - L^2 - C - L^3 - R^2}_{\xi} = \underbrace{\frac{\xi}{\xi} - L^2}_{\xi} - \underbrace{\frac{R^2}{\xi} - L^2}_{\xi}$$

In re Application of: Khan, Saeed R. PATENT
Application No: 10/596,751 Attorney Docket No. JHU6010-1
Filled: October 18, 2007

Page 3

(2)

(3) wherein

n is 0 or 1:

 X^{2} is -N(R⁴)- or -CH(R⁴)-:

 L^2 , L^3 , L^4 , and L^5 are is independently a bond, substituted or unsubstituted alkylene, substituted or unsubstituted heteroalkylene, or substituted or unsubstituted heterocycloalkylene; and

R², R³, and R⁴ are independently hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted exploalkyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl.

wherein if X is C=, L^2 is a bond, L^3 is unsubstituted alkylene, R^1 is (1), and R^3 is para to L^4 -B(OH)₂, then R^2 is not substituted or unsubstituted aryl or substituted or unsubstituted heteroaryl.

2. (Currently Amended) The compound of claim 1, wherein

In re Application of: Khan, Saeed R. Application No.: 10/596,751

Filed: October 18, 2007

Page 4

PATENT

Attorney Docket No. JHU6010-1

 $\rm L^{1}$ is a bond, substituted or unsubstituted $\rm C_{1}\text{-}C_{20}$ alkylene, or substituted or unsubstituted 2 to 20 membered heteroalkylene;

 L^2 , L^3 , L^4 , and L^5 are is independently a bond, substituted or unsubstituted $C_{1^-}C_{20}$ alkylene, substituted or unsubstituted 2 to 20 membered heteroalkylene, or substituted or unsubstituted 3 to 8 membered heterocycloalkylene; and

 R^2 , R^3 , and R^4 are independently hydrogen, substituted or unsubstituted C_1 - C_{20} alkyl, substituted or unsubstituted 2 to 20 membered heteroalkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, substituted or unsubstituted 3 to 8 membered heterocycloalkyl, substituted or unsubstituted or unsubstituted heteroaryl.

3. (Currently Amended) The compound of claim 1, wherein

 L^1 is a bond, unsubstituted C_1 - C_{10} alkylene, or unsubstituted 2 to 10 membered heteroalkylene;

 L^2 , L^3 , L^4 -and L^5 -are <u>is</u> independently a bond, unsubstituted C_1 - C_{10} alkylene, unsubstituted 2 to 10 membered heteroalkylene, or unsubstituted 3 to 8 membered heterocycloalkylene; and

 R^2 , R^3 , and R^4 are independently hydrogen, substituted or unsubstituted C_1 - C_{10} alkyl, substituted or unsubstituted 2 to 10 membered heteroalkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl, substituted or unsubstituted 3 to 8 membered heterocycloalkyl, substituted or unsubstituted or unsubstituted heteroaryl.

4. (Currently Amended) The compound of claim 1, wherein

 L^1 is abond a bond, substituted or unsubstituted C_1 - C_{10} alkylene, or substituted or unsubstituted 2 to 10 membered heteroalkylene: and

In re Application of: Khan, Saeed R. PATENT
Application No.: 10/596,751 Attorney Docket No. JHU6010-1

Filed: October 18, 2007

Page 5

 L^2 , L^3 , L^4 -and L^5 are \underline{is} independently a bond, substituted or unsubstituted C_1 - C_{10} alkylene, substituted or unsubstituted 2 to 10 membered heteroalkylene, or substituted or unsubstituted 3 to 8 membered heterocycloalkylene;

5. (Currently Amended) The compound of claim 1, wherein R¹ has the formula (1), wherein

L² is a bond, unsubstituted C₁-C₁₀ alkylene, unsubstituted 2 to 10 membered heteroalkylene, or unsubstituted 3 to 8 membered heterocycloalkylene;

L³ is a bond, unsubstituted C₄·C₁₀ alkylene, unsubstituted 2 to 10 membered heteroalkylene, or unsubstituted 3 to 8 membered heterocycloalkylene; and

 R^2 is R^{21} -substituted or unsubstituted C_{1} - C_{15} alkyl, R^{21} -substituted or unsubstituted 2 to 10 membered heteroalkyl, R^{21} -substituted or unsubstituted C_{3} - C_{8} cycloalkyl, R^{21} -substituted or unsubstituted 3 to 8 membered heterocycloalkyl, R^{21} -substituted or unsubstituted aryl, or R^{21} -substituted or unsubstituted beteroaryl, wherein

 R^{21} is a halogen, -OH, -SH, -NH₂, -CF₃, -B(OH)₂, -C(0)NHOH, unsubstituted $C_1\text{-}C_{10}$ alkyl, unsubstituted 2 to 10 membered heteroalkyl, unsubstituted $C_3\text{-}C_8$ cycloalkyl, unsubstituted 3 to 8 membered heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, or -OR 22 , wherein

R²² is unsubstituted C₃-C₁₀ alkyl, unsubstituted 2 to 10 membered heteroalkyl, unsubstituted C₃-C₈ cycloalkyl, unsubstituted 3 to 8 membered heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, or -(CH₂)_qB(OH)₂, wherein q is an integer from 1 to 5.

(Currently Amended) The compound of claim 1, wherein R¹ has the formula (I), wherein
 L² is a bond, or unsubstituted 3 to 8 membered heterocycloalkylene:

L3 is unsubstituted C1-C10 alkylene; and

In re Application of: Khan, Saeed R. Application No.: 10/596,751 Filed: October 18, 2007

Page 6

 R^2 is R^{21} -substituted or unsubstituted C_1 - C_{15} alkyl, or R^{21} -substituted or unsubstituted aryl, wherein

 R^{21} is halogen, -OH, -SH, -NH₂, -CF₃, -B(OH)₂, -C(0)NHOH, unsubstituted C_1 -C $_{10}$ alkyl, unsubstituted 2 to 10 membered heteroalkyl, unsubstituted C_3 -C $_8$ cycloalkyl, unsubstituted 3 to 8 membered heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, or -OR 22 , wherein

R²² is unsubstituted C₁-C₁₀ alkyl, or unsubstituted 2 to 10 membered heteroalkyl.

7. (Currently Amended) The compound of claim 6, wherein

L2 is a bond or piperazinylene;

L3 is unsubstituted C1-C3 alkenylene;

R² is unsubstituted C₁-C₁₅ alkyl, or R²¹-substituted or unsubstituted aryl, wherein

 R^{21} is halogen, -C(0)NHOH, unsubstituted C_1 - C_{10} alkyl, unsubstituted 2 to 10 membered heteroalkyl, or -OR²², wherein

R²² is unsubstituted C₁-C₁₀ alkyl, or unsubstituted 2 to 10 membered heteroalkyl.

 (Currently Amended) The compound of claim 1, wherein R¹ has the formula (2) or (3), wherein

L⁴ and L⁵ are independently a bond, unsubstituted C₄-C₁₀ alkylene, unsubstituted 2 to 10 membered heteroalkylene, or unsubstituted 3 to 8 membered heterocycloalkylene;

 R^4 is unsubstituted C_1 - C_{10} alkyl, unsubstituted 2 to 10 membered heteroalkyl, unsubstituted C_3 - C_8 cycloalkyl, unsubstituted 3 to 8 membered heterocycloalkyl, unsubstituted aryl, or unsubstituted heteroaryl;

In re Application of: Khan, Saeed R. Application No.: 10/596,751 Filed: October 18, 2007 PATENT Attorney Docket No. JHU6010-1

Page 7

 R^3 is R^{31} -substituted or unsubstituted C_{1} - C_{10} alkyl, R^{31} -substituted or unsubstituted 2 to 10 membered heteroalkyl, R -substituted or unsubstituted C_{3} - C_{8} cycloalkyl, R^{31} -substituted or unsubstituted 3 to 8 membered heterocycloalkyl, R^{31} -substituted or unsubstituted aryl, or R^{31} -substituted or unsubstituted heteroaryl, wherein

 R^{31} is halogen, -OH, -SH, -NH₂, -CF₃, -B(OH)₂, -C(0)NHOH, unsubstituted C_1 - C_{10} alkyl, unsubstituted 2 to 10 membered heteroalkyl, unsubstituted C_3 - C_8 cycloalkyl, unsubstituted 3 to 8 membered heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, or -OR³², wherein

R³² is unsubstituted C₁-C₁₀ alkyl, unsubstituted 2 to 10 membered heteroalkyl, unsubstituted C₃-C₈ cycloalkyl, unsubstituted 3 to 8 membered heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, or -(CH2)_mB(OH)₂, wherein m is an integer from 1 to 5.

 (Currently Amended) The compound of claim 1, wherein R¹ has the formula (1) or (2), wherein

L4 and L5 are unsubstituted C1-C10 alkylene;

R4 is unsubstituted C1-C10 alkyl;

R3 is R31-substituted arvl, wherein

 R^{31} is a halogen, -C(0)NHOH, unsubstituted C_1 - C_{10} alkyl, unsubstituted 2 to 10 membered heteroalkyl, or -OR³², wherein

 R^{32} is an unsubstituted C_1 - C_{10} alkyl, unsubstituted 2 to 10 membered heteroalkyl, or -(CH₂)_mB(OH)₂, wherein m is 1 to 5.

(Canceled).

- Page 8
- 11. (Original) The compound of claim 1, wherein R¹ is para to -L¹-B(OH)₂.
- 12. (Original) The compound of claim 1, wherein X is -C=.
- 13. (Original) The compound of claim 1, wherein L¹ is a bond or methylene.
- 14. (Withdrawn) A method of treating a tumor or cancer in a patient in need thereof comprising administering to said patient an effective amount of a compound having the Formula:

wherein

X1 is -O or -N=:

 ${\bf L}^1$ is a bond, substituted or unsubstituted alkylene, or substituted or unsubstituted heteroalkylene; and

R1 has the formula:

(1)
$$\begin{cases} \frac{1}{2} - \frac{1}{2} - \frac{1}{2} - \frac{1}{2} - \frac{1}{2} \\ 0 \end{cases}$$
, (2) $\begin{cases} \frac{1}{2} - \frac{1}{2} - \frac{1}{2} - \frac{1}{2} \\ \frac{1}{2} - \frac{1}{$

In re Application of: Khan, Saeed R. Application No.: 10/596,751 Filed: October 18, 2007 Page 9

wherein

n is 0 or 1:

 X^{2} is -N(R⁴)- or -CH(R⁴)-:

L², L³, L⁴, and L⁵ are independently a bond, substituted or unsubstituted alkylene, substituted or unsubstituted heteroalkylene, or substituted or unsubstituted heterocycloalkylene; and

 R^2 , R^3 , and R^4 are independently hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl.

wherein if X is -C=, L^2 is a bond, L^3 is unsubstituted alkylene, R^1 is (1), and R^1 is para to $-L^1$ -B(OH)₂, then R^2 is not substituted or unsubstituted aryl or substituted or unsubstituted heteroaryl.

15. (Withdrawn) The method of claim 14, wherein said tumor is selected from the group consisting of breast, cervical, stomach, colon, bladder, rectal, liver, pancreatic, lung, cervix uteri, corpus uteri, ovary, prostate, testis, renal, brain/ons, head, neck, throat, anal and oral cancers, eye or ocular cancer, skin melanoma, Ewing's Sarcoma, Kaposi's Sarcoma, basal cell carcinoma and squamous cell carcinoma, small cell lung cancer, mouth/pharynx, esophageal, larynx, kidney and lymphoma, acute lymphocytic leukemia, and acute myelogenous leukemia.

Page 10

16. (Withdrawn) A method of inhibiting MDM2 expression in a mammal, comprising administering an amount of a compound effective to inhibit said expression, said compound having the Formula:

wherein

X1 is -O or -N=:

 \mathcal{L}^1 is a bond, substituted or unsubstituted alkylene, or substituted or unsubstituted heteroalkylene; and

R1 has the formula:

(2)
$$R^4 / n$$
 , or

In re Application of: Khan, Saeed R. Application No.: 10/596,751 Filed: October 18, 2007 Page 11

wherein

n is 0 or 1;

 X^{2} is -N(R⁴)- or -CH(R⁴)-:

L², L³, L⁴, and L⁵ are independently a bond, substituted or unsubstituted alkylene, substituted or unsubstituted heteroalkylene, or substituted or unsubstituted heterocycloalkylene; and

R², R³, and R⁴ are independently hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl.

wherein if X is -C=, L^2 is a bond, L^3 is unsubstituted alkylene, R^1 is (1), and R^1 is para to $-L^1$ -B(OH)₂, then R^2 is not substituted or unsubstituted aryl or substituted or unsubstituted heteroaryl.

17. (New) The compound of claim 1, wherein R¹ has the formula:

$$(2) \qquad \qquad \begin{pmatrix} \chi_1 \\ \chi_2 \\ R^4 \end{pmatrix}_n \qquad \text{or (3)}$$

18. (New) The compound of claim 1, wherein the compound having the formula:

HO B L1

has formulae:

$$(HO)_{z}B \longrightarrow HO)_{z}$$

wherein R' is halogen, alkyl, heteroalkyl, alkoxy, heteroalkoxy, or hydroxamic acid.

19. (New) A compound having formula:

Page 13

wherein

 X^1 is C= or -N=:

 \mathbf{L}^1 is a bond, substituted or unsubstituted alkylene, or substituted or unsubstituted heteroalkylene;

R1 has the formula:

(3) wherein

(2)

n is 0 or 1;

In re Application of: Khan, Saeed R. Application No.: 10/596,751 Filed: October 18, 2007 Page 14

PATENT Attorney Docket No. JHU6010-1

 R^2 , R^3 , and R^4 are independently hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted eycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl.